


Order CYPRINIFORMES

Cigar Barbs

Family LEPTOBARBIDAE

Bleeker 1864



Possibly first-published image of *Leptobarbus hoevenii*. From: Bleeker, P. 1863–64. Atlas ichthyologique des Indes Orientales Néerlandaises, publié sous les auspices du Gouvernement colonial néerlandais. Tome III. Cyprins. 1–150, Pls. 102–144. [This plate published 1863.]

Leptobarbus

Bleeker 1859

leptós (Gr. λεπτός), thin, referring to slender shape of *L. hoevenii*;
Barbus (Cyprinidae: Barbinae), its original genus

***Leptobarbus hoevenii* (Bleeker 1851)** in honor of Dutch physician-zoologist Jan van der Hoeven (1801–1868), “le célèbre professeur de zoologie”

***Leptobarbus hosii* (Regan 1906)** in honor of Charles Hose (1863–1929), British colonial administrator in Borneo, zoologist and ethnologist, who collected holotype

***Leptobarbus melanopterus* Weber & de Beaufort 1916** black-finned, from *mélanos* (Gr. μέλανος), genitive of *mélas* (μέλας), black, and *pterus*, from *pterón* (Gr. πτερόν) or *ptéryx* (πτέρυξ), wing or fin, referring to nearly all-black ventral, pectoral and dorsal fins

***Leptobarbus melanotaenia* Boulenger 1894** *mélanos* (Gr. μέλανος), genitive of *mélas* (μέλας), black; *taenia*, from *tainía* (Gr. ταινία), band or ribbon, referring to black lateral stripe running along body above lateral line and/or black vertical bar on posterior border of gill opening

***Leptobarbus rubripinna* (Fowler 1937)** *ruber* (L.), red; *pinna* (L.) fin, referring to “more or less bright orange red” coloration of fins (except dorsal) in living specimens