# The ETYFish Project

COMMENTS: **=** v. 4.0 - 17 Sept. 2023

# **Order MUGILIFORMES**

#### Family MUGILIDAE Mullets

25 genera · 79 species

#### Agonostomus Bennett 1832

a-, without; gonos, angle; stomus, mouth, referring to more rounded (less angular) lower jaw compared to Mugil

#### Agonostomus catalai Pellegrin 1932

in honor of René Catala (1901-1988), coffee planter and amateur (at the time) biologist in Madagascar, who collected holotype

# Agonostomus telfairii Bennett 1832

in honor of Irish botanist and ship surgeon Charles Telfair (1778-1833), who collected and/or acquired plants and animals in Mauritius in 1810, including type of this mullet

## Aldrichetta Whitley 1945

-etta, diminutive connoting endearment: in honor of Fred C. Aldrich, Chief Inspector of Fisheries and Game (Perth, Western Australia) from 1911-1937, an "able and enthusiastic fisheries investigator"

#### Aldrichetta forsteri (Valenciennes 1836)

in honor of Johann Reinhold Forster (1729-1798), naturalist aboard James Cook's second voyage on HMS *Resolution*, who observed this mullet in New Zealand in 1769; while he identified it as *Mugil albula* (=cephalus), his notes provided sufficient descriptive data for Valenciennes to determine it was a separate and new species

## Cestraeus Valenciennes 1836

latinization of kestreus, a name used by Aristotle for mullets

# Cestraeus goldiei (Macleay 1883)

in honor of "Alex" Goldie, probably Andrew Goldie (1840-1891), Scottish-born merchant, explorer and natural history collector, who collected type from the Goldie River (which he named after himself) in Papua New Guinea

#### Cestraeus oxyrhyncus Valenciennes 1836

oxy, sharp; rhynchus, snout, referring to sharp angle of lower lip, which makes the snout more pointed

## Cestraeus plicatilis Valenciennes 1836

pleated, referring to its "very thick" upper lip, with a "fleshy mass" at the end of the snout, "covered with excessively fine and numerous papillae, which give this organ a velvety appearance" (translations)

## Chaenomugil Gill 1863

chaeno-, gape, "distinguished by the longitudinal cleft of the mouth"; mugil, mullet

#### Chaenomugil proboscideus (Günther 1861)

having a proboscis, referring to "extremely thick, conically produced" thick part of upper lip

#### Chelon Artedi 1793

ancient Greek name mentioned by Aristotle as a synonym for *kephalos* (see *Mugil cephalus*, below) [authorship sometimes attributed to Röse, who reprinted Artedi's 1738 publication]

# Chelon auratus (Risso 1810)

gilded, referring to the "beautiful golden spots that adorn its operculum" (translation)

#### Chelon bandialensis (Diouf 1991)

-ensis, suffix denoting place: Bandiala, one of three coastal rivers that constitute the Sine Saloum estuary in Senegal, type locality

# Chelon bispinosus (Bowdich 1825)


bi-, two; spinosus, thorny, allusion not explained, perhaps referring to two spines (plus seven rays) in second dorsal fin (first dorsal fin consists of four "strong" spines with no rays)

#### Chelon caeruleus Deef 2018

blue, referring to dark-blue color of sides [originally spelled *caeruleum* but emended to *caeruleus* since *Chelon* is masculine; of uncertain taxonomic status, provisionally included here]

#### Chelon dumerili (Steindachner 1870)

in honor of August Duméril (1812-1870), herpetologist and ichthyologist, Muséum national d'Histoire naturelle



Chelon luciae. From: Penrith, M.-L. and M. J. Penrith. 1967. A new species of mullet of the genus Ellochelon (Pisces: Mugilidae) from St. Lucia Estuary, Zululand. Durban Museum Novitates v. 8 (pt 8): 69-75.

(Paris), who loaned Steindachner specimens of other Sengalese fishes for study

#### Chelon labrosus (Risso 1827)

thick-lipped, allusion not explained but probably referring to its thicker lips compared to C. ramada

# Chelon luciae (Penrith & Penrith 1967)

of St. Lucia estuary, Kwazulu-Natal, South Africa, type locality

# Chelon natalensis (Castelnau 1861)

-ensis, suffix denoting place: KwaZulu-Natal, South Africa, locality (also occurs in Swaziland and Mozambique)

#### Chelon parsia (Hamilton 1822)

local name for this mullet in West Bengal, India

# Chelon persicus Senou, Randall & Okiyama 1995

Persian, referring to Persian Gulf, where it appears to be endemic

#### Chelon ramada (Risso 1827)

from ramado, local name for this mullet along the northwestern Mediterranean Sea

# Chelon richardsonii (Smith 1846)

patronym not identified, probably in honor of surgeon-naturalist John Richardson (1787-1865)

# Chelon saliens (Risso 1810)

leaping or jumping, referring to how this mullet "jumps with extraordinary velocity when it finds himself trapped in a net" (translation)

#### Chelon tricuspidens (Smith 1935)

referring to its relatively large tricuspid teeth

#### Crenimugil Schultz 1946

creni-, referring to crenulate lips of C. crenilabis; mugil, mullet

# Crenimugil buchanani (Bleeker 1853)

in honor of Francis Hamilton-Buchanan (1762-1829), Scottish physician and naturalist, whose 1822 account of Indo-Gangetic fishes (including descriptions of three mullets) is cited several times by Bleeker

#### Crenimugil crenilabis (Forsskål 1775)

creni-, crenulate; labis, lip, referring to row of crenulate papillae along external edge of its thick lips

# Crenimugil heterocheilos (Bleeker 1855)

heteros, different; cheilos, lip, allusion not explained, presumably referring to lip papillae, different from most other mullet genera

# Crenimugil seheli (Forsskål 1775)

Sehêli, Arabic name for this mullet

# Crenimugil pedaraki (Valenciennes 1836)

from *Peddaraki Sover*e, its local name at Vizagapatam on the Coromandel Coast of India, as reported by Russell (1803)

#### Daiaus Valenciennes 1836

latinization of dajao, local name for this mullet in Puerto Rico

## Dajaus monticola (Bancroft 1834)

of the mountains, referring to its occurrence in steep forested streams as high as 1500 m

#### Ellochelon Whitley 1930

Ellon Gistel 1848, an unneeded and presumably forgotten replacement name for Mugil, which for some reason Whitley wished to commemorate; chelon, ancient Greek name mentioned by Aristotle as a synonym for kephalus (see Mugil cephalus)

#### Ellochelon vaigiensis (Quoy & Gaimard 1825)

-ensis, suffix denoting place: Pulau Waigeo (or Vaigiou), Papua Barat, Indonesia, type locality (but widely occurs in Red-Sea and Indo-West Pacific)

# **Gracilimugil Whitley 1941**

gracilis, slender or thin, referring to its compressed body and "graceful" form; mugil, mullet

#### Gracilimugil argenteus (Quoy & Gaimard 1825)

silvery, referring to the silvery coloration of its "fine and pearly" scales (translation)

#### Joturus Poey 1860

latinization of Joturo, Cuban name for J. pichardi in Havana (also known as Foturo)

#### Joturus pichardi Poey 1860

in honor of Cuban lexicographer-geographer Esteban Pichardo (1799-1879), "estimable auteur" of *Diccionario Provincial de voces Cubanos* and *Geografia de la isla de Cuba* (he also recommended the vernacular name *Joturo*)

#### Mugil Linnaeus 1758

Latin for mullet, possibly derived from mulgeo, to suck, referring to how M. cephalus feeds by sucking up sediment

#### Mugil bananensis (Pellegrin 1927)

-ensis, suffix denoting place: Banana, Democratic Republic of the Congo, type locality (occurs along eastern Atlantic coast from Senegal to Angola, and Cape Verde Island)

#### Mugil brevirostris Miranda Ribeiro 1915

brevis, short; rostris, snout, referring to its "de focinho curto"

# Mugil broussonnetii Valenciennes 1836

in honor of physician-naturalist Pierre Marie Auguste Broussonet (1761-1807), who acquired type from Joseph Banks, collected during James Cook's first great voyage (1768-1771)

# Mugil capurrii (Perugia 1892)

in honor of Capt. Guiseppe Capurro, who collected type in Senegal, and to whom the Museo Civico di Genova "owes many interesting animals collected by him during his travels" (translation)

#### Mugil cephalus Linnaeus 1758

kephalos, a name dating to Aristotle (it is not clear if the name relates to cephalus, meaning head)

# Mugil curema Valenciennes 1836

Curema, Portuguese vernacular used by Dutch naturalist Jorge Marcgrave in his 1648 Historia Naturalis Brasiliae, doubtless corresponding to the Spanish vernacular querimana (or queriman)

#### Mugil curvidens Valenciennes 1836

curvus, bent; dens, teeth, referring to teeth in the lower jaw, which are directed downwards and forwards

#### Mugil hospes Jordan & Culver 1895

a landlord, referring to the near constant presence in its mouth or branchial cavity of a small oniscid crustacean similar to that found in the Menhaden, *Brevoortia tyrannus* (Clupeidae)

# Mugil hubensis Rizvi & Iqbal 1986

-ensis, suffix denoting place: Hub River, Baluchistan, Pakistan, type locality

#### Mugil incilis Hancock 1830

pertaining to a ditch or trench, "being chiefly found in the trenches or ditches dug for draining the flat lands of the coast of Guiana"

#### Mugil liza Valenciennes 1836

vernacular name for "mullet with big lips" in Spain, apparently applied to this western Atlantic and Caribbean species by the Spanish and/or Valenciennes

## Mugil longicauda Guitart Manday & Alvarez-Lajonchere 1976

longus, long; cauda, tail, referring to its "notably elongated caudal peduncle"

#### Mugil margaritae Menezes, Nirchio, Oliveira & Siccharamirez 2015

of Isla Margarita, Venezuela, type locality

#### Mugil rubrioculus Harrison, Nirchio, Oliveira, Ron & Gaviria 2007

rubri-, red; oculus, eye, referring to its distinctly reddish-orange iris

#### Mugil setosus Gilbert 1892

bristly, referring to "much larger multiserial setæ" (hair-like structures) on premaxilla compared to the similar *M. curema* (perhaps referring to a sympatric population later described as *M. hospes*)

## Mugil thoburni (Jordan & Starks 1896)

in honor of Wilbur Wilson Thoburn (1859-1899), who taught bionomics (ecology) at Stanford University (where Jordan was president), for his work on sculpins (Cottidae)

## Mugil trichodon Poey 1875

trichos, hair; odon, tooth, presumably referring to the filiform appearance of its teeth

#### Myxus Günther 1861

mucous or slime, allusion not explained, perhaps referring to how mullets often feed on "soft organic substances"

### Myxus elongatus Günther 1861

elongate, referring to how its "body appears to be more elongate" than other mullet species

#### Neochelon Durand, Chen, Shen, Fu & Borsa 2012

neo-, new, i.e., a new genus of Chelon

#### Neochelon falcipinnis (Valenciennes 1836)

falcatus, sickle-shaped; pinnis, fin, referring to falcate anal fin

# Neomyxus Steindachner 1878

neo-, new, proposed as a subgenus of Myxus, i.e., a new Myxus

## Neomyxus leuciscus (Günther 1871)

*leukiskos*, Greek word for chub (probably derived from *leukos*, white), allusion not explained, perhaps referring to its silvery sides and white belly and/or its resemblance to the European cyprinid *Leuciscus leuciscus* 

#### Oedalechilus Fowler 1903

oedaleus, swollen; cheilus, lip, referring to fleshy upper lip

#### Oedalechilus labeo (Cuvier 1829)

labeo, one with large lips, referring to its fleshy upper lip, thicker than Mugil chelo (=Chelon labrosus), its presumed congener at the time

# Osteomugil Luther 1982


osteo-, bone, referring to how genus is differentiated by an osteological character (absent vs. present posterior zygapophysial hooks on any trunk vertebra); mugil, mullet

# Osteomugil cunnesius (Valenciennes 1836)

latinization of *Kunnesee*, its local name at Vizagapatam on the Coromandel Coast of India, as reported by Russell (1803)

# Osteomugil engeli (Bleeker 1858)

per Bleeker (1860), in honor of Chris Engel, who provided many of the illustrations in Bleeker's Atlas Ichthyologique des Orientales Neerlandaises



Neomyxus leuciscus. From: Günther, A. 1872. Report on several collections of fishes recently obtained for the British Museum. Proceedings of the Zoological Society of London 1871 (pt 3) (art. 1): 652-675, Pls. 53-70.

## Osteomugil perusii (Valenciennes 1836)

in honor of French naval officer and explorer Jean-François de Galaup La Pérouse (1741-1788), "a name that will remind all scientists of the coasts where this species can be found" (translation), referring to Vanikoro Island in the South Pacific (type locality), where Pérouse was stranded after both his ships struck reefs; he and his crew were never seen again

#### Osteomugil robustus (Günther 1861)

stout, presumably referring to its "very thick" body

#### Osteomugil speigleri (Bleeker 1858)

patronym not identified but almost certainly in honor of Ludwig Speigler (1824-1893), who provided many of the illustrations in Bleeker's *Atlas Ichthyologique des Orientales Neerlandaises* 

#### Parachelon Durand, Chen, Shen, Fu & Borsa 2012

para-, near, allusion not explained, presumably referring to its placement next to Chelon on the authors' phylogenetic tree

## Parachelon grandisquamis (Valenciennes 1836)

grandis, large; squamus, scale, referring to larger scales, 26 or 28 along lateral line, compared to 45 on Mugil cryptocheilos (=Chelon auratus) and Mugil plumieri (=cephalus), its presumed congeners at the time

#### Paramugil Ghasemzadeh, Ivantsoff & Aarn 2004

para-, near; Mugil, genus in which both included species had previously been placed

#### Paramugil georgii (Ogilby 1897)

etymology not explained, perhaps referring to the Georges River, New South Wales, Australia, type locality

#### Paramugil parmatus (Cantor 1849)

shielded, allusion not explained, perhaps referring to large silvery-white scales covering (i.e., shielding) body

#### Planiliza Whitley 1945

planus, flat, presumably referring to "broad, flat and depressed" interorbital of *P. ordensis*; Liza (=Chelon), its presumed closest relative (see Mugil liza for etymology)

## Planiliza abu (Heckel 1843)

from Abu Sukkanejn, Arabic name for this mullet in Mosul, Iraq, from Abu, father (i.e., possessor, or one with), and sukkanejn, anchor, probably referring to its toothed suborbital bone (per Heckel), or abu, a prefix particle denoting "an animal having or being like" and sukkanejn, dual steering wheel or handle bars, referring to two dorsal (i.e., steering) fins (per Mikaili & Shayegh 2011, "An etymological review on fish common and scientific names in the Euphrates and Tigris")

# Planiliza alata (Steindachner 1892)

winged, presumably referring to prolonged and sickle-shaped ventral, anal, caudal and second dorsal fins

#### Planiliza carinata (Valenciennes 1836)

keeled, referring to predorsal scales in midline thickened and ridged to form a keel

# Planiliza haematocheila (Temminck & Schlegel 1845)

blood-lipped, from *hematos*, bloody, and *cheilus*, lip, referring to "dirty purple" (translation) of lips of living specimens [often spelled as a nondeclinable noun, *hematocheilus*]

## Planiliza klunzingeri (Day 1888)

in honor of German physician and zoologist Carl Benjamin Klunzinger (1834-1914), who pointed out that Day had incorrectly identified this mullet as *Mugil carinatus* (=*P. carinata*) in 1876

# Planiliza lauvergnii (Eydoux & Souleyet 1850)

patronym not identified, probably in honor of Barthélemy Lauvergne (1805-1871), painter and draughtsman on voyage that collected type

# Planiliza macrolepis (Smith 1846)

macro-, large; lepis, scale, referring to "very large and hard" scales of back and sides

# Planiliza mandapamensis (Thomson 1997)

-ensis, suffix denoting place: Kilakarei, south of Mandapam, southern India, type locality

#### Planiliza melinoptera (Valenciennes 1836)

melinus, quince-yellow; pterus, fin, referring to lemon-yellow fins of living specimens, especially the caudal fin (based on manuscript description and illustration by Quoy & Gaimard)

# Planiliza ordensis (Whitley 1945)

-ensis, suffix denoting place: Ord River, Western Australia, type locality (also occurs in New Guinea and Tonga)

## Planiliza planiceps (Valenciennes 1836)

planus, flat; ceps, head, referring to its "extremely flattened" (translation) head

#### Planiliza subviridis (Valenciennes 1836)

sub-, less than or somewhat; viridis, green, referring to its greenish-gray back

## Planiliza tade (Forsskål 1775)

Tâde, Arabic name for this mullet

# Plicomugil Schultz 1953

plicatus, folded, referring to its folded and lobed lips; mugil, mullet

#### Plicomugil labiosus (Valenciennes 1836)

large-lipped, referring to its fleshy lips

# Pseudomyxus Durand, Chen, Shen, Fu & Borsa 2012

pseudo-, false, i.e., although this genus may resemble Myxus, such an appearance is false

## Pseudomyxus capensis (Valenciennes 1836)

-ensis, suffix denoting place: Cape of Good Hope, South Africa, type locality

#### Rhinomugil Gill 1863

rhino-, snout, "distinguished by [its] projecting rounded snout"; mugil, mullet

#### Rhinomugil corsula (Hamilton 1822)

latinization of Khorsula, local name for this mullet in West Bengal, India

# Sicamugil Fowler 1939

sica, dagger, referring to strongly spinate preorbital spine; mugil, mullet

## Sicamugil cascasia (Hamilton 1822)

presumably a local Gangetic name for this mullet [sometimes placed in *Minimugil* Senou 1988, proposed in an unpublished Ph.D. dissertation and therefore not available; authorship sometimes dated to Durand 2016, who cited Senou and provided some diagnostic traits of the genus, but name is not "explicitly indicated as intentionally new" (ICZN 16.1) and therefore remains unavailable]

## Sicamugil hamiltonii (Day 1870)

in honor of Francis Hamilton-Buchanan (1762-1829), Scottish physician and naturalist, who described the similar *S. cascasia* in 1822

#### Squalomugil Ogilby 1908

squalus, shark, referring to flattened head and inferior mouth; mugil, mullet

# Squalomugil nasutus (De Vis 1883)

large-nosed, referring to its "rounded, sharp, [and] much produced" snout [sometimes placed in Rhinomugil]

# Trachystoma Ogilby 1888

trachys, rough; stoma, mouth, referring to vomer and palate furnished with distinct bands of villiform teeth (but jaws are toothless)

#### Trachystoma petardi (Castelnau 1875)

in honor of Mr. Petard (forename not available), who sent fishes to Castelnau collected from the Richmond River, New South Wales, including type of this one